

IBS

IRRITABLE BOWEL SYNDROME

This is one of the most common conditions affecting the gut. Individuals usually complain of recurrent abdominal pain which may vary in site and severity and is often associated with variable stool form and frequency. Abdominal distension or bloating may be a feature. IBS is a functional condition; the symptoms are absolutely genuine but are not due to an identifiable disease process or detectable abnormality on investigation.

Studies suggest that IBS is much more common in women. The symptoms may follow an infection or an episode of food poisoning in about 20% of cases (so-called post-infective IBS). IBS may be worse at times of stress. Physical examination, stool culture and blood test results are always normal in IBS; if an abnormality is detected then the diagnosis should be reconsidered.

Treatment is difficult but usually involves treating the predominant symptom. For example, if diarrhoea is the main problem then loperamide may be tried. In people with post-infective, diarrhoea-predominant IBS ondansetron or colesevalam (in case of co-existing bile salt malabsorption) may be of some help.

There is a new treatment for people with moderate-severe IBS with constipation (IBS-C). The drug is called linaclotide and may be prescribed to improve symptoms of abdominal pain, bloating and constipation in adults with IBS-C. See Patient Information sheet for more information.

Treating pain may be more challenging and a variety of different drugs may be tried; simple remedies such as peppermint oil capsules,

alverine or mebeverine may be tried first and if unsuccessful, progressing to hyoscine (Buscopan). In more severe cases, drugs affecting the levels of serotonin in the gut (and the brain), such as citalopram, may be tried or in those whom are unkeen on drugs, cognitive therapy or hypnotherapy may be considered.

Some people with diarrhoea and abdominal bloating thought to be due to IBS may respond to treatment with probiotics, antibiotics (see Patient Information sheet on Small Intestine Bacterial Overgrowth) or a gluten-free diet (in absence of coeliac disease).

There is increasing interest in the relationship between certain dietary sugars and IBS. Researchers from Monash University in Australia found that removing certain fermentable sugars from the diet helped up to 75% of people with IBS. A low FODMAP (Fermentable, Oligosaccharides, Disaccharides, Monosaccharides And Polyols) diet. These substances may be poorly absorbed in the small intestine and then act as a food source for bacteria in the large intestine (colon); it is this process which may produce gas and draws in liquid leading to nausea, flatulence, diarrhoea and bloating.

For more information on IBS or any of the treatment options please contact West Kent Gastroenterology (phone: 01892 526726; email info@westkentgastroenterology.com).

Written by Dr Adam Harris
Consultant Gastroenterologist